


DOVEKIE

Alle alle alle**Regional Occurrence** (BCRs 14, 30)

- Distribution: pelagic in region
- Habitats used: mostly pelagic, along seacoasts (non-breeding); crevices on steep coastal cliffs (breeding)

NAWCP BD=5; NAWCP ND=2

Regional Abundance

- 1980s population size: 10,000-100,000nb
- Historical population size: unknown
- % change in regional population: unknown

NAWCP PS=4; NAWCP PT=3

Regional Threats

- Mortality can be due to rough seas and/or failing plankton supply
- Degradation/loss of habitat at inland nesting areas

NAWCP TB=2; NAWCP TN=3

Conservation Status: Abundance (b=breeding, nb=nonbreeding, t=total individuals) and Conservation Rank

Global	North America	BCR 14	BCR 30	Mid-Atlantic	S. New England	Gulf of Maine	Maritimes
16,000,000- 36,000,000t	1,000b	?nb	?nb	?nb	?nb	?nb	?nb
IUCN- Not Evaluated	NAWCP- Moderate Concern (PI<1%)	Moderate (PI=?)	Moderate (PI=?)	NJ- SZN; DE- no info; MD-SZN; VA-SN	NY-S?N; CT- no info; RI- SZN MA-SN; VT- no info	MA-SN; NH-SZN ME-S1N; NB- S4N, NS-S5N	QC-SN; NB- S4N PE-SZN; NS-S5N

Conservation Needs: *Sustainability:* 77% hatched chicks fledged; 35% females to rear surviving fledgling; *Commuting Range:* 2.5-100 km, may shorten for breeding birds during chick rearing; generally < 17 km; dives to depth of 35-50m; cold waters in moderate to heavy offshore pack ice, over banks at sea and at upwellings and oceanographic fronts that transport and concentrate planktonic prey near the surface; feed along ice edges; *Issues:* ; *Population Goals:* ; *Habitat Goals:*